

ADDITION: Saturday, December 19, 16:00 - 16:40

A semilinear elliptic problem with a singularity at $u = 0$

FRANÇOIS MURAT

Laboratoire Jacques-Louis Lions
Université Pierre et Marie Curie (Paris VI) and CNRS
Paris, France

email: murat@ann.jussieu.fr

In this joint work with Daniela Giachetti (Rome, Italy) and Pedro J. Martínez Aparicio (Cartagena, Spain) (see [3] and [4]), we consider the semilinear elliptic equation with homogeneous Dirichlet boundary condition

$$-div A(x)Du = F(x, u) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega, \quad u \geq 0 \quad \text{in } \Omega,$$

where the nonlinearity $F(x, u)$ is singular at $u = 0$, and more precisely where F is a Carathéodory function $F : \Omega \times [0, +\infty[\rightarrow [0, +\infty]$ which satisfies

$$0 \leq F(x, s) \leq \frac{h(x)}{\Gamma(s)} \quad \text{a.e. } x \in \Omega, \forall s > 0,$$

with $h \geq 0$, $h \in L^r(\Omega) \subset H^{-1}(\Omega)$ and $\Gamma : [0, +\infty[\rightarrow [0, +\infty[$ a C^1 , Lipschitz-continuous, nondecreasing function such that $\Gamma(0) = 0$ and $\Gamma(s) > 0$ for every $s > 0$. A model for such a function $F(x, s)$ is for example given by

$$F(x, s) = \frac{f(x)}{\exp(-\frac{1}{s})} \left(2 + \sin\left(\frac{1}{s}\right) \right) + \frac{g(x)}{s^\gamma} + l(x) \quad \text{a.e. } x \in \Omega, \forall s > 0,$$

where the functions f , g and l are nonnegative and belong to $L^r(\Omega)$.

The main difficulty is to give a convenient definition of the solution of this problem, in particular when $\Gamma(s) \ll s$ for s close to 0.

We give such a definition and we prove the existence and stability of this solution, as well as its uniqueness when $F(x, s)$ is non increasing in s .

This work has been inspired by the papers [2] of Lucio Boccardo and Luigi Orsina and [1] of Lucio Boccardo and Juan Casado-Diaz.

References:

- [1] L. Boccardo & J. Casado-Diaz: Some properties of solutions of some semilinear elliptic singular problems and applications to the G-convergence, *Asymptotic Analysis*, 86 (2104), 1–15.
- [2] L. Boccardo & L. Orsina: Semilinear elliptic equations with singular nonlinearities, *Calculus of Variations and Partial Differential Equations*, 37 (2010), 363–380.
- [3] D. Giachetti, P.J. Martínez-Aparicio & F. Murat: An elliptic equation with a mild singularity at $u = 0$: existence and homogenization, to appear.
- [4] D. Giachetti, P.J. Martínez-Aparicio & F. Murat: Definition, existence, stability and uniqueness of the solution to a semilinear elliptic problem with a strong singularity at $u = 0$, to appear.

Program

17th December		18th December		19th December
8:30-9:00 Registration				
9:00-9:40 Opening		9:00-9:40 Piatnitski		9:00-9:40 Kinderlehrer
9:45-10:25 Tartar		9:45-10:25 Buttazzo		9:45-10:25 Conca
10:30-10:50 Coffee break		10:30-10:50 Coffee break		10:30-10:50 Coffee break
10:50-11:30 Raoult		10:50-11:30 Cioranescu		10:50-11:30 Sequeira
11:35-12:15 Seppecher		11:35-12:15 Gaudiello		11:35-12:15 Donato
12:20-13:00 Fragalà		12:20-13:00 Oliveira		12:20-13:00 Bismut
13:05-14:30 Lunch		13:05-14:30 Lunch		13:05-14:30 Lunch
14:30-15:10 Bouchitté		14:30-15:10 Dias		14:30-15:10 Leoni
15:15-15:55 Allaire		15:15-15:55 Beirão da Veiga		15:15-15:55 Sanchez
16:00-16:40 Francfort		16:00-16:40 Rodrigues		16:00-16:40 Murat
16:45-17:15 Coffee break		16:45-17:15 Coffee break		16:45-17:45 Coffee and Poster session
Section A	Section B	Section A	Section B	
17:15-17:30 Toader	17:15-17:30 Santos	17:15-17:30 Zappale	17:15-17:30 Monsaingeon	
17:35-17:50 Giannetti	17:35-17:50 Quítalo	17:35-17:50 Almi	17:35-17:50 Antonic	
17:55-18:10 Morandotti	17:55-18:10 Coelho	17:55-18:10 Burazin	17:55-18:10 Kreisbeck	17:50-18:30 Fonseca
18:15-18:30 Barbarosie	18:15-18:30 Tavares	18:15-18:30 Vorotnikov	18:15-18:30 Velcic	18:35-19:00 Closing
		20:00-23:00 Dinner		